

Exercises to the Lecture FSVT

Prof. Dr. Klaus Madlener

sheet 12

**Exercise 1:** [standard combinators]

Prove the following equations to be valid for the standard combinators  $I \equiv \lambda x.x$ ,  $K \equiv \lambda xy.x$ ,  $B \equiv \lambda xyz.x(yz)$ ,  $K_* \equiv \lambda xy.y$ ,  $S \equiv \lambda xyz.xz(yz)$ :

1.  $IM = M$ , 2.  $KMN = M$ , 3.  $K_*MN = N$ , 4.  $SMNL = ML(NL)$ , 5.  $BLMN = L(M(N))$

**Exercise 2:** [Number presentations]

Let the following number presentations in the  $\lambda$ -calculus be given:

1.  $c_0 \equiv \lambda fx.x$ ,  $c_{n+1} \equiv \lambda fx.f^{n+1}(x)$
2.  $d_0 \equiv I$ ,  $d_{n+1} \equiv [\text{false}, d_n]$
3.  $z_0 \equiv KI$ ,  $z_{n+1} \equiv SBz_n$ ,

where  $F^0(M) \equiv M$ ,  $F^{n+1}(M) \equiv F(F^n(M))$ ,  $\text{true} \equiv K$ ,  $\text{false} \equiv K_*$ ,  $[M, N] \equiv \lambda z.zMN$ .


Prove:

1. There are terms  $T, T^{-1}$  with  $Tc_n \equiv d_n$  and  $T^{-1}d_n \equiv c_n$  for all  $n$ .
2. There are terms  $R, R^{-1}$  with  $Rd_n \equiv z_n$  and  $R^{-1}z_n \equiv d_n$  for all  $n$ .

**Exercise 3:** [properties of redexes]

1. Make yourself familiar with the notation used in chapter 11 of the lecture. Use the following paper: Bergstra, Klop :: Conditional Rewrite Rules: Confluence and Termination. JCSS 32 (1986)
2. Prove the following Lemma (Lemma 11.5 on slide 361).

Let  $D$  be an elementary reduction's diagram for orthogonal systems,  $R_i \subseteq M_i$  ( $i = 0, 2, 3$ ) redexes with  $R_0 \dots \rightarrow R_2 \dots \rightarrow R_3$  i.e  $R_2$  is Rest of  $R_0$  and  $R_3$  is Rest of  $R_2$ . Then there is a unique redex  $R_1 \subseteq M_1$  with  $R_0 \dots \rightarrow R_1 \dots \rightarrow R_3$ , i.e.



**Delivery: until 31.01.2011,**  
**by E-Mail to huechting@informatik.uni-kl.de**